

Tabelle 2

Variabel	Level	n	Failures	Univariate Cox-Regression				Multiple Cox-Regression			
				HR	95 %	CI	Pval	HR	95 %	CI	Pval
Alter bei Therapiebeginn	<65	164	19	Ref.			<b>0,03</b>				
	65–69	149	19	1,18	0,62	2,22					
	70–74	162	14	0,79	0,40	1,57					
	75+	143	23	<b>2,02</b>	1,09	3,71					
Prostatavolumen (ml)	<27	130	18	1,03	0,55	1,95	0,86				
	27–34,9	150	21	1,10	0,59	2,02					
	35–43,9	131	15	0,82	0,42	1,61					
	44+	156	20	Ref.							
Initialer PSA (ng/ml)	< 10	460	45	Ref.			<b>&lt;0,0001</b>				
	10–20	102	18	<b>2,76</b>	1,59	4,79					
	20+	48	12	<b>3,72</b>	1,96	7,05					
Gleason-Score	<6	37	2	0,39	0,09	1,67	<b>&lt;0,0001</b>	0,39	0,09	1,67	<b>&lt;0,0001</b>
	6	211	22	1,08	0,59	1,96		1,08	0,59	1,96	
	7a	269	21	Ref.				Ref.			
	7b	66	13	<b>2,70</b>	1,35	5,40		<b>2,70</b>	1,35	5,40	
	8	9	3	<b>5,80</b>	1,72	19,5		<b>5,80</b>	1,72	19,5	
	9–10	25	14	<b>12,5</b>	6,28	24,7		<b>12,5</b>	6,28	24,7	
Gleason-Gruppierung nach ISUP2014 / WHO2016	1	248	24	0,94	0,52	1,69	<b>&lt;0,0001</b>				
	2	269	21	Ref.							
	3	66	13	<b>2,70</b>	1,35	5,39					
	4	9	3	<b>5,77</b>	1,71	19,4					
	5	25	14	<b>12,4</b>	6,25	24,6					
D'Amico-Gruppierung	Low	211	21	0,75	0,43	1,30					
	Intermediär	331	32	Ref.							
	High	70	22	<b>4,17</b>	2,42	7,18	<b>&lt;0,0001</b>				
Anteil positiver Biopsien	<0,1	77	8	1,56	0,65	3,17	<b>0,0005</b>				
	0,1–0,19	123	11	1,00	0,46	2,18					
	0,2–0,39	156	15	Ref.							
	0,4+	136	28	<b>3,19</b>	1,69	6,03					
Befall der Seitenlappen	Einseitig	296	35	Ref.			<b>0,28</b>				
	Beidseitig	195	27	1,32	0,80	2,18					

Tab. 2: Patientencharakteristika sowie Hazard Ratios des biochemischen rezidivfreien Überlebens (bRFS).